

Table 3E
2004 Pregnancy Nutrition Surveillance ⁽¹⁾
North Dakota
Comparison of Racial and Ethnic Distribution by Region

Page: 1
Run Date: 1 / 18 / 06

Comparison of Racial and Ethnic Distribution ⁽²⁾							
Region	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other/Unknown %
CDHU	114	79.8	0.0	0.9	19.3	0.0	0.0
FDHU	509	79.6	5.5	4.3	7.9	2.8	0.0
James R	224	92.9	1.8	2.2	3.1	0.0	0.0
LRDHU	194	47.4	0.0	1.0	51.0	0.5	0.0
North RR	546	74.9	2.4	11.9	9.2	1.6	0.0
Rolette	236	6.4	0.0	0.0	93.6	0.0	0.0
So Central	352	78.4	1.1	1.1	19.0	0.3	0.0
South RR	701	81.9	6.0	4.6	5.8	1.7	0.0
SWDHU	187	94.1	0.0	2.1	2.7	1.1	0.0
UMDHU	161	90.1	0.6	0.6	8.7	0.0	0.0
Unknown Regions	18	*	*	*	*	*	*
North Dakota	3,242	74.2	2.8	4.3	17.5	1.2	0.0
Nation (Prior Year)	747,174	48.7	24.4	21.2	1.3	1.9	2.5

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 4E
2004 Pregnancy Nutrition Surveillance ⁽¹⁾
North Dakota
Comparison of Age Distribution by Region

Comparison of Age Distribution ⁽²⁾								
Region	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
CDHU	114	0.0	4.4	9.6	55.3	28.1	2.6	0.0
FDHU	509	0.0	3.9	12.4	69.5	13.2	0.6	0.4
James R	224	0.0	4.0	12.1	62.5	19.6	0.0	1.8
LRDHU	194	0.5	9.3	12.4	62.9	13.9	1.0	0.0
North RR	546	0.0	5.1	11.7	62.1	19.6	1.3	0.2
Rolette	236	1.7	7.6	19.1	56.8	13.6	1.3	0.0
So Central	352	0.6	4.5	13.9	64.8	14.5	1.1	0.6
South RR	701	0.1	2.1	12.8	68.8	15.4	0.4	0.3
SWDHU	187	0.0	1.6	12.3	71.1	13.9	0.5	0.5
UMDHU	161	0.0	3.7	17.4	59.6	18.0	1.2	0.0
Unknown Regions	18	*	*	*	*	*	*	*
North Dakota	3,242	0.2	4.3	13.2	64.8	16.1	0.9	0.4
Nation (Prior Year)	747,174	0.4	6.9	13.4	60.1	17.9	1.3	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 5E
2004 Pregnancy Nutrition Surveillance ⁽¹⁾
North Dakota
Comparison of Maternal Health Indicators by Region

Comparison of Maternal Health Indicators (2, 3)												
Region	Prepregnancy BMI (4)				Weight Gain (5)				Anemia (Low Hb/Hct) (6)			
	Under-weight		Over-weight	< Ideal		> Ideal		3rd Trimester		Postpartum		
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)		
CDHU	109	10.1 (7)	47.7 (8)	88	* *	* *	12	* *	62	* *		
FDHU	486	8.8 (3)	41.6 (1)	345	14.8 (6)	47.2 (4)	75	* *	311	26.4 (3)		
James R	217	6.9 (1)	53.0 (10)	144	7.6 (1)	50.7 (5)	25	* *	136	19.1 (1)		
LRDHU	189	7.9 (2)	46.6 (6)	137	16.8 (8)	58.4 (9)	30	* *	121	30.6 (5)		
North RR	506	10.5 (8)	43.5 (3)	265	13.6 (5)	43.8 (1)	62	* *	290	34.8 (6)		
Rolette	235	8.9 (4)	50.2 (9)	189	9.0 (4)	58.2 (8)	23	* *	150	24.7 (2)		
So Central	337	11.9 (9)	47.2 (7)	259	15.8 (7)	52.1 (6)	50	* *	217	36.4 (7)		
South RR	651	13.1 (10)	42.2 (2)	485	19.0 (9)	45.6 (3)	99	* *	397	38.3 (8)		
SWDHU	170	9.4 (5)	43.5 (3)	142	8.5 (3)	56.3 (7)	26	* *	105	27.6 (4)		
UMDHU	152	9.9 (6)	46.1 (5)	103	7.8 (2)	44.7 (2)	14	* *	84	* *		
Unknown Regions	16	* *	* *	10	* *	* *	3	* *	13	* *		
North Dakota	3,068	10.3	45.0	2,167	14.4	49.5	419	24.6	1,886	30.4		
Nation (Prior Year)	678,999	12.1	43.0	579,271	25.2	44.2	99,365	30.3	253,236	29.5		

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one region's rate to other regions. Rank 1 = best rate.

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 6E
2004 Pregnancy Nutrition Surveillance ⁽¹⁾
North Dakota
Comparison of Maternal Behavioral Indicators by Region

Comparison of Maternal Behavioral Indicators ^(2, 3)						
Region	Medical Care				WIC Enrollment	
	1st Trimester		No Care		1st Trimester	
	Number	% (Rank)	% (Rank)		Number	% (Rank)
CDHU	103	90.3 (3)	1.0 (3)		91	* *
FDHU	434	86.6 (6)	0.5 (1)		483	36.2 (9)
James R	192	90.6 (2)	2.1 (5)		197	44.7 (5)
LRDHU	175	73.1 (10)	4.0 (8)		186	38.7 (6)
North RR	452	92.3 (1)	2.2 (6)		470	45.5 (3)
Rolette	213	78.9 (9)	1.9 (4)		234	51.3 (1)
So Central	311	88.7 (5)	2.3 (7)		299	46.8 (2)
South RR	570	90.0 (4)	0.9 (2)		554	45.3 (4)
SWDHU	173	83.2 (7)	4.6 (10)		170	36.5 (8)
UMDHU	143	81.1 (8)	4.2 (9)		136	36.8 (7)
Unknown Regions	16	* *	* *		14	* *
North Dakota	2,782	87.0	1.9		2,834	42.7
Nation (Prior Year)	440,121	77.1	8.9		695,912	27.6

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one region's rate to other regions. Rank 1 = best rate.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 7E
2004 Pregnancy Nutrition Surveillance (1)
North Dakota
Comparison of Smoking Indicators by Region

Comparison of Smoking Indicators (2, 3)										
Region	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)		Number	% (Rank)	Number	% (Rank)	
CDHU	106	50.0 (7)	99	* *	33	* *	73	* *	92	* *
FDHU	414	42.3 (1)	386	20.2 (2)	121	43.8 (2)	393	31.3 (3)	350	27.4 (4)
James R	198	49.5 (6)	165	26.7 (7)	60	* *	165	31.5 (4)	160	28.1 (5)
LRDHU	188	55.3 (9)	151	29.8 (8)	62	* *	167	32.9 (6)	148	26.4 (3)
North RR	430	44.2 (3)	306	18.0 (1)	93	* *	401	27.2 (1)	302	20.9 (1)
Rolette	235	63.8 (10)	205	45.4 (9)	123	19.5 (4)	218	44.5 (9)	205	23.4 (2)
So Central	316	50.6 (8)	298	24.5 (5)	119	45.4 (1)	271	36.9 (8)	287	35.5 (9)
South RR	676	44.7 (4)	551	23.0 (3)	183	32.2 (3)	558	32.4 (5)	546	30.6 (8)
SWDHU	168	44.0 (2)	160	24.4 (4)	59	* *	146	28.8 (2)	157	29.3 (6)
UMDHU	155	47.1 (5)	119	25.2 (6)	36	* *	107	33.6 (7)	118	29.7 (7)
Unknown Regions	15	* *	12	* *	5	* *	12	* *	11	* *
North Dakota	2,901	47.7	2,452	25.1	894	36.5	2,511	32.8	2,376	28.1
Nation (Prior Year)	617,683	29.1	483,916	18.9	121,114	31.9	294,627	25.5	331,615	22.3

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one region's rate to other regions. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 8E
2004 Pregnancy Nutrition Surveillance ⁽¹⁾
North Dakota
Comparison of Infant Health Indicators by Region

Comparison of Infant Health Indicators (2, 3)														
Region	Birthweight				Preterm (6)	Full Term LBW (7)		Ever Breastfed						
	Low (4)		High (5)											
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)				
CDHU	88	*	*	*	*	67	*	*	53	*	*	93	*	*
FDHU	383	3.9	(1)	11.0	(7)	375	9.1	(1)	319	1.6	(3)	408	61.8	(1)
James R	172	8.7	(9)	8.1	(1)	151	13.2	(7)	126	2.4	(6)	176	50.6	(6)
LRDHU	154	3.9	(1)	14.3	(9)	152	10.5	(2)	132	0.8	(1)	159	37.7	(8)
North RR	329	6.7	(7)	9.7	(4)	287	11.5	(5)	232	2.2	(4)	351	58.7	(4)
Rolette	111	5.4	(3)	10.8	(6)	108	11.1	(3)	96	*	*	107	33.6	(9)
So Central	287	5.6	(6)	8.4	(2)	247	11.3	(4)	214	2.3	(5)	295	48.8	(7)
South RR	542	5.5	(4)	10.0	(5)	413	12.3	(6)	360	1.4	(2)	547	58.5	(5)
SWDHU	163	5.5	(4)	9.2	(3)	146	13.7	(8)	125	2.4	(6)	165	61.2	(3)
UMDHU	118	6.8	(8)	11.0	(7)	91	*	*	75	*	*	120	61.7	(2)
Unknown Regions	15	*	*	*	*	12	*	*	11	*	*	15	*	*
North Dakota	2,362	5.6		10.1		2,049	11.8		1,743	1.8		2,436	54.9	
Nation (Prior Year)	616,163	8.0		7.5		599,892	11.3		502,591	3.8		616,123	55.7	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one region's rate to other regions. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 3E
2002 - 2004 Pregnancy Nutrition Surveillance ⁽¹⁾
North Dakota
Comparison of Racial and Ethnic Distribution by Region
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾							
Region	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other/Unknown %
CDHU	395	85.3	0.3	1.3	12.9	0.3	0.0
FDHU	1,671	81.3	5.1	4.2	7.3	2.0	0.0
James R	716	94.1	1.0	2.8	2.0	0.1	0.0
LRDHU	718	50.0	0.0	0.6	49.2	0.3	0.0
North RR	1,703	75.0	2.7	12.5	8.0	1.8	0.0
Rolette	786	6.5	0.0	0.0	93.3	0.3	0.0
So Central	1,105	78.6	0.7	1.7	18.4	0.5	0.0
South RR	2,331	82.2	5.9	4.6	6.0	1.3	0.0
SWDHU	529	93.0	0.2	1.9	4.0	0.9	0.0
UMDHU	483	83.6	0.6	1.7	14.1	0.0	0.0
Unknown Regions	101	77.2	1.0	14.9	5.9	1.0	0.0
North Dakota	10,538	74.2	2.8	4.5	17.5	1.1	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 4E
2002 - 2004 Pregnancy Nutrition Surveillance ⁽¹⁾
North Dakota
Comparison of Age Distribution by Region
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Region	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
CDHU	395	0.0	5.1	10.4	61.3	21.3	1.5	0.5
FDHU	1,671	0.1	3.5	13.6	69.1	12.3	1.0	0.4
James R	716	0.0	4.9	12.3	61.2	20.1	0.7	0.8
LRDHU	718	0.1	7.8	14.9	60.2	15.5	1.3	0.3
North RR	1,703	0.0	4.8	13.2	65.0	15.7	1.0	0.3
Rolette	786	1.0	9.2	17.6	57.4	14.1	0.6	0.1
So Central	1,105	0.3	4.0	16.7	61.8	15.3	1.3	0.7
South RR	2,331	0.2	3.7	13.1	67.1	14.9	0.7	0.2
SWDHU	529	0.2	3.0	14.6	67.3	14.0	0.8	0.2
UMDHU	483	0.0	5.0	20.1	58.0	15.3	1.4	0.2
Unknown Regions	101	2.0	5.0	22.8	57.4	11.9	1.0	0.0
North Dakota	10,538	0.2	4.7	14.4	64.2	15.2	0.9	0.4

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 5E
2002 - 2004 Pregnancy Nutrition Surveillance ⁽¹⁾
North Dakota
Comparison of Maternal Health Indicators by Region
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)										
Region	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)			
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)
CDHU	379	12.1 (7)	42.2 (5)	300	17.0 (9)	48.0 (5)	51	* *	222	24.8 (5)
FDHU	1,589	9.9 (2)	41.2 (3)	1,150	13.8 (5)	50.5 (6)	248	17.3 (1)	1,008	25.7 (7)
James R	696	9.5 (1)	49.0 (9)	487	12.3 (3)	45.4 (2)	66	* *	446	20.6 (3)
LRDHU	700	10.3 (4)	46.3 (7)	464	17.9 (10)	51.7 (8)	115	27.0 (3)	418	23.2 (4)
North RR	1,568	12.1 (7)	41.9 (4)	921	14.4 (6)	46.1 (3)	202	23.3 (2)	871	31.0 (8)
Rolette	779	10.0 (3)	49.4 (10)	600	10.8 (1)	55.3 (10)	94	* *	555	19.8 (2)
So Central	1,049	10.5 (6)	45.7 (6)	792	14.6 (7)	51.3 (7)	152	30.9 (5)	660	35.3 (9)
South RR	2,195	12.8 (9)	39.8 (1)	1,645	16.6 (8)	46.6 (4)	334	27.2 (4)	1,356	36.4 (10)
SWDHU	481	12.9 (10)	41.0 (2)	357	12.3 (3)	52.1 (9)	70	* *	263	25.1 (6)
UMDHU	441	10.4 (5)	46.9 (8)	245	11.0 (2)	44.5 (1)	35	* *	186	16.1 (1)
Unknown Regions	97	* *	* *	64	* *	* *	11	* *	68	* *
North Dakota	9,974	11.2	43.3	7,025	14.5	49.0	1,378	23.6	6,053	28.3

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one region's rate to other regions. Rank 1 = best rate.

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 6E
2002 - 2004 Pregnancy Nutrition Surveillance ⁽¹⁾
North Dakota
Comparison of Maternal Behavioral Indicators by Region
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)						
Region	Medical Care				WIC Enrollment	
	1st Trimester		No Care		1st Trimester	
	Number	% (Rank)	% (Rank)		Number	% (Rank)
CDHU	351	90.0 (2)	0.6 (1)		347	31.1 (10)
FDHU	1,435	85.5 (6)	1.3 (3)		1,553	36.0 (8)
James R	632	87.5 (3)	2.5 (6)		628	46.7 (2)
LRDHU	597	74.5 (10)	2.5 (6)		694	38.3 (7)
North RR	1,397	87.3 (4)	3.5 (8)		1,436	44.3 (3)
Rolette	709	77.9 (9)	1.3 (3)		768	49.7 (1)
So Central	949	87.0 (5)	1.9 (5)		931	42.0 (5)
South RR	1,935	90.1 (1)	0.6 (1)		1,900	44.3 (3)
SWDHU	475	82.7 (7)	4.6 (9)		465	39.4 (6)
UMDHU	431	79.4 (8)	5.6 (10)		393	31.8 (9)
Unknown Regions	90	* *	* *		91	* *
North Dakota	9,001	85.5	2.1		9,206	41.6

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one region's rate to other regions. Rank 1 = best rate.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 7E
2002 - 2004 Pregnancy Nutrition Surveillance ⁽¹⁾
North Dakota
Comparison of Smoking Indicators by Region
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Region	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
CDHU	360	52.8 (7)	343	31.2 (9)	132	35.6 (5)	283	36.0 (7)	332	32.2 (9)
FDHU	1,352	43.9 (3)	1,286	23.2 (3)	420	36.0 (4)	1,274	31.4 (3)	1,223	27.0 (3)
James R	655	44.1 (4)	560	25.2 (5)	187	30.5 (8)	539	31.5 (4)	548	24.1 (2)
LRDHU	701	52.9 (8)	520	29.2 (7)	211	35.5 (6)	615	33.8 (6)	512	27.9 (4)
North RR	1,388	43.7 (2)	1,079	18.3 (1)	335	44.5 (1)	1,272	27.9 (1)	1,069	21.9 (1)
Rolette	774	66.8 (10)	661	46.1 (10)	414	22.2 (9)	730	42.6 (10)	660	29.7 (7)
So Central	1,008	52.6 (6)	922	26.7 (6)	368	38.6 (3)	847	37.5 (8)	890	31.2 (8)
South RR	2,223	44.2 (5)	1,880	25.0 (4)	615	33.0 (7)	1,836	31.9 (5)	1,867	29.2 (6)
SWDHU	476	43.1 (1)	411	20.2 (2)	136	42.6 (2)	404	28.7 (2)	405	28.4 (5)
UMDHU	465	54.4 (9)	305	30.5 (8)	92	* *	254	39.4 (9)	302	34.8 (10)
Unknown Regions	90	* *	72	* *	35	* *	81	* *	70	* *
North Dakota	9,492	48.3	8,039	26.3	2,945	34.5	8,135	33.1	7,878	28.1

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one region's rate to other regions. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 8E
2002 - 2004 Pregnancy Nutrition Surveillance ⁽¹⁾
North Dakota
Comparison of Infant Health Indicators by Region
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
Region	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
CDHU	318	5.0 (1)	6.9 (1)	270	11.1 (4)	232	1.3 (1)	326	49.7 (8)
FDHU	1,277	6.0 (4)	9.9 (6)	1,197	11.4 (6)	1,013	2.3 (4)	1,337	60.1 (2)
James R	582	8.8 (10)	7.9 (3)	513	10.7 (1)	444	3.2 (9)	599	52.1 (7)
LRDHU	535	6.7 (8)	13.8 (10)	532	11.7 (8)	450	2.9 (8)	556	42.3 (9)
North RR	1,098	6.1 (5)	11.0 (7)	913	11.1 (4)	776	1.7 (3)	1,133	58.9 (3)
Rolette	459	5.7 (3)	12.6 (9)	426	10.8 (2)	376	2.7 (6)	460	30.4 (10)
So Central	896	8.0 (9)	7.7 (2)	715	13.4 (10)	608	3.3 (10)	911	55.4 (6)
South RR	1,837	6.6 (6)	9.3 (5)	1,432	10.8 (2)	1,269	2.3 (4)	1,849	57.7 (4)
SWDHU	427	5.6 (2)	8.4 (4)	378	11.4 (6)	321	2.8 (7)	444	56.8 (5)
UMDHU	303	6.6 (6)	11.6 (8)	210	12.4 (9)	183	1.6 (2)	307	63.2 (1)
Unknown Regions	83	* *	* *	75	* *	67	* *	86	* *
North Dakota	7,815	6.6	9.8	6,661	11.3	5,739	2.4	8,008	54.8

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one region's rate to other regions. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.